

# Molecular Gas in the Host Galaxy of a Quasar at Redshift $z=6.42$

Fabian Walter<sup>1\*</sup>, Frank Bertoldi<sup>+</sup>, Chris Carilli<sup>\*</sup>, Pierre Cox<sup>++</sup>, K.Y. Lo<sup>\*</sup>,  
Roberto Neri<sup>&</sup>, Xiaohui Fan<sup>\*\*</sup>, Alain Omont<sup>#</sup>, Michael A. Strauss<sup>##</sup>, Karl  
M. Menten<sup>+</sup>

\* National Radio Astronomy Observatory, P.O. Box 0, Socorro, NM 87801, USA

+ Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

++ Institut d'Astrophysique Spatiale, Universite de Paris-Sud, 91405 Orsay, France

& IRAM, 300 Rue de la Piscine, 38406 St-Martin-d'Heres, France

\*\* Steward Observatory, University of Arizona, 933 N. Cherry Ave., Tucson, AZ 85721, USA

# Institut d'Astrophysique de Paris, CNRS & Université Paris 6, 75014 Paris, France

## Princeton University Observatory, Princeton, NJ 08544, USA

<sup>1</sup> Jansky Fellow

Revised Nature manuscript: 2003-05-04710

Observations of the molecular gas phase in quasar host galaxies provide fundamental constraints on galaxy evolution at the highest redshifts. Molecular gas is the material out of which stars form; it can be traced by spectral line emission of carbon–monoxide (CO). To date, CO emission was detected in more than a dozen quasar host galaxies with redshifts ( $z$ ) larger 2, the record holder being at  $z=4.69^{1-3}$ . At these distances the CO lines are shifted to longer wavelengths, enabling their observation with sensitive radio and millimetre interferometers<sup>4</sup>. Here we present the discovery of CO emission toward the quasar SDSS J114816.64+525150.3 (hereafter J1148+5251) at a redshift of  $z=6.42^{5,6}$  (when the universe was only  $\sim 1/16$  of its present age). This is the first detection of molecular gas at the end of cosmic reionization. The presence of large amounts of molecular gas ( $M(\text{H}_2)=2.2\times 10^{10} M_\odot$ ) in an object at this time demonstrates that heavy element enriched molecular gas can be generated rapidly in the earliest galaxies.

The source J1148+5251 is a luminous quasar which is thought to be powered by mass accretion onto a supermassive black hole of mass  $1\text{--}5\times 10^9 M_\odot^7$ . Optical spectra of J1148+5251 show a clear Gunn–Peterson trough<sup>5,6</sup> (i.e. Ly  $\alpha$  absorption by the neutral intergalactic medium), indicating that this quasar is situated at the end of the epoch of reionization<sup>8</sup>. Thermal emission from warm dust was detected at millimetre wavelengths, implying a far infrared (FIR) luminosity of  $1.3 \times 10^{13} L_\odot^9$ , corresponding to about 10% of the bolometric luminosity of the system (we assume  $H_0=71 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_\Lambda=0.73$  and  $\Omega_m=0.27$  throughout). We searched for molecular gas in J1148+5251 using the National Radio Astronomy Observatory’s (NRAO) Very Large Array (VLA) and the IRAM Plateau de Bure interferometer (PdBI) in February–April 2003. The VLA observations have higher spatial resolution,

whereas the PdBI has better spectral resolution. We observed the CO(3-2), (6-5) and (7-6) lines which are shifted to 46.6, 93.2 and 108.7 GHz, respectively, at  $z \sim 6.42$ . For J1148+5251 the possible redshift range based on broad optical emission lines is  $z = 6.35^5$  to  $6.43^7$ . We searched this entire range for CO(3-2) emission using the VLA at a resolution of 50 MHz ( $320 \text{ km s}^{-1}$ ,  $\Delta z = 0.008$ ) per channel. CO(3-2) emission is clearly detected at high Signal to Noise in the channel centred at 46.6149 GHz (see Figs. 1 and 2), corresponding to a redshift of  $z = 6.418 \pm 0.004$ . The PdBI covered a redshift range from  $z = 6.40$ – $6.44$  and the CO emission is also detected at high significance (Fig. 2)<sup>10</sup>. In this letter we concentrate on the VLA results; a more detailed analysis of the entire dataset will be presented elsewhere<sup>10</sup>.

The velocity-integrated CO(3-2) flux is  $S_{\text{CO}(3-2)} \Delta v = 0.18 \pm 0.04 \text{ Jy km s}^{-1}$ . On the Kelvin scale ( $1 \text{ Jy} = 250 \text{ K}$  for a  $1.5''$  beam) the source has a peak line flux of  $0.14 \text{ K}$  and a line integral of  $44 \pm 10 \text{ K km s}^{-1}$ . The CO luminosity is given by<sup>11</sup>:

$$L' = 3.25 \times 10^7 \times S_{\text{CO}} \Delta v [\text{Jy km s}^{-1}] \times \nu_{\text{obs}}^{-2} [\text{GHz}] \times D_L^2 [\text{Mpc}] \times (1 + z)^{-3} \text{ K km s}^{-1} \text{ pc}^2,$$

where  $D_L$  is the luminosity distance. For J1148+5251 we obtain:  $L'_{\text{CO}(3-2)} = 2.7 \times 10^{10} \text{ K km s}^{-1} \text{ pc}^2$ .

The total molecular gas mass,  $M(\text{H}_2)$ , can be derived from the relation  $M(\text{H}_2) = \alpha \times L'_{\text{CO}(1-0)}$ .

In Galactic molecular clouds the conversion factor is  $\alpha \sim 2.0 \text{ M}_\odot (\text{K km s}^{-1} \text{ pc}^2)^{-1}$ <sup>12</sup>, but here we use the value  $\alpha \sim 0.8 \text{ M}_\odot (\text{K km s}^{-1} \text{ pc}^2)^{-1}$ , appropriate for ultraluminous infrared galaxies<sup>13</sup> (ULIRGs,  $L_{\text{FIR}} \geq 10^{12} L_\odot$ ) and nuclear starburst galaxies<sup>14</sup>. Assuming a constant brightness temperature from CO(3-2) to CO(1-0)<sup>10</sup>, this leads to  $M(\text{H}_2) = 2.2 \times 10^{10} \text{ M}_\odot$  for J1148+5251.

If the FIR luminosity of J1148+5251 is powered by star formation, the implied star formation rate (SFR) is  $3000 \text{ M}_\odot \text{ yr}^{-1}$ <sup>9</sup>. Comparing this to the molecular gas mass implies

a short gas depletion timescale of order  $10^7$  years. However, given the high luminosity of the quasar it is possible that some fraction of the dust is heated by the quasar, in which case the star formation rate quoted above becomes an upper limit (resulting in a longer gas depletion timescale). The FIR continuum-to-CO line ratio for J1148+5251 is large,  $440 L_{\odot} (\text{K km s}^{-1} \text{pc}^2)^{-1}$ , which is an order of magnitude larger than for normal star forming galaxies  $(5-50)^{15}$ , but is comparable to that seen in ULIRGs and other high redshift quasars<sup>9</sup>. Some authors have suggested that high continuum-to-line ratios indicate high star formation efficiencies<sup>15</sup>. Alternatively, dust heating by the AGN could lead to large continuum-to-line ratios.

A dynamical mass for the system can be derived from the CO observations, assuming that the CO is rotationally supported. The upper limit to the CO source diameter derived from Gaussian fitting is  $1.5''$  ( $1''=5.46 \text{ kpc}$ ). A lower limit to the source diameter can be derived from the measured brightness temperature ( $T_{\text{obs}}$ ) by assuming an intrinsic brightness temperature  $T_{\text{b}}$ :  $\Omega_S/\Omega_B=(T_{\text{obs}}/T_{\text{b}})(1+z)$ , where  $\Omega_S$  and  $\Omega_B$  are the source and beam solid angles. The most extreme case is a single optically thick emission region, in which case  $T_{\text{b}}$  equals  $T_{\text{ex}}-T_{\text{CMB}}$ , the excitation brightness temperature minus the continuum temperature of the cosmic background at  $z=6.4$  ( $\sim 20 \text{ K}$ ). This gives a minimum source diameter of  $0.2''(T_{\text{b}}/50 \text{ K})^{-1/2}$ . Assuming a rotational velocity of  $v_{\text{rot}} = 130 \sin^{-1}(i) \text{ km s}^{-1}$  (Fig. 2), where  $i$  is the inclination angle with respect to the sky plane, yields a range for the dynamical mass of  $M_{\text{dyn}} = 2 \text{ to } 16 \times 10^9 \times \sin^{-2}(i) M_{\odot}$ . This is comparable to the molecular gas mass in J1148+5251, implying that either molecular gas dominates the dynamics of the system, or that the plane of rotation is close to the sky plane. The optical spectrum of the quasar shows no significant reddening<sup>5</sup>, despite the large dust mass of the host galaxy<sup>9</sup>. This suggests that the disk of molecular gas (and dust) is oriented close to the sky plane such that the line-of-sight to the nucleus is relatively unobscured. We note that the mass of the central black

hole<sup>4</sup> likely constitutes a significant fraction ( $\sim 1 - 10\%$ ) of the total dynamical mass in the inner few kpc of J1148+5251, in contrast to what is generally found for nearby galaxies<sup>16,17</sup>.

The CO redshift of  $z=6.419$  (Fig. 2) corresponds to the systemic redshift of the host galaxy of J1148+5251, since it traces the extended molecular gas distribution in the quasar and is not associated with emission supposedly emerging from energetic processes, e.g. lines of shocked outflow gas or gas related to AGN accretion. The latter has been found to be shifted significantly in frequency with respect to the systemic redshift<sup>18</sup>. Indeed, studies of high ionization UV lines (in particular SiIV)<sup>5</sup> in J1148+5251 yield a redshift of  $z=6.37$  corresponding to a velocity offset of  $\sim 2000 \text{ km s}^{-1}$ . The CO redshift is however in good agreement with the redshift derived from the low ionization MgII line<sup>7</sup>.

Measuring an accurate redshift is particularly important to determine the state of the IGM around the quasar (the ‘proximity effect’). In the optical spectrum<sup>5,7</sup> there is essentially zero emission present in the wavelength range corresponding to Ly $\alpha$  at  $z=5.7-6.33$  (due to the Gunn–Peterson effect); emission for  $z > 6.33$  can be attributed to the ionized medium around the quasar. Using the source redshift of 6.419 results in a Strömgen sphere around J1148+5251 with a comoving radius of  $R_S=4.8 \text{ Mpc}$ . An estimate for the age for this sphere (and hence for the quasar itself) can be derived from  $R_S$  using<sup>19,6</sup>

$$\dot{N}_{ph}/(10^{58} \text{ s}^{-1}) \times t_q/(10^7 \text{ yr}) = 0.34 \times [R_S/(4.8 \text{ Mpc}) \times (1 + z_q)/7.419]^3$$

and assuming  $\dot{N}_{ph} = 0.2 - 1.3 \times 10^{58} \text{ s}^{-1}$ <sup>20,21,6</sup>. This calculation results in an age of order  $10^7 \text{ yr}$  for the quasar activity in J1148+5251. Interestingly, this timescale is comparable to the formation timescale for the central black hole, which has an e-folding timescale for accretion of order  $4 \times 10^7 \text{ yr}$  (assuming a radiative efficiency of 0.1 and that the black hole

is accreting at the Eddington limit)<sup>7</sup>, suggesting that the AGN ionizes a significant volume around the quasar.

The fact that we detect CO emission implies that the process of enrichment of the ISM in J1148+5251 with heavy elements is relatively advanced, i.e., that significant star formation has occurred in the quasar host galaxy prior to  $z=6.4$ . This conclusion is supported by the fact that both strong metal emission lines<sup>5</sup>, and thermal emission from warm dust<sup>9</sup>, were detected from J1148+5251. A recent optical study of a  $z=6.28$  quasar even suggests supersolar metallicities in these early objects<sup>22</sup>. Assuming a Galactic abundance for J1148+5251 we derive an order of magnitude estimate for the total mass in CO of  $M(\text{CO}) \sim 3 \times 10^7 M_\odot$ <sup>10</sup>. This amount of C and O could be produced relatively quickly by  $\sim 10^7$  hundred solar mass Population III stars<sup>23</sup> with lifetimes  $< 10^7$  yr. Likewise, if enrichment was achieved by conventional supernovae<sup>24</sup> and assuming a star formation rate of order  $3000 M_\odot \text{yr}^{-1}$ <sup>9</sup>, we estimate a time for enrichment of order  $10^7$  yr. We consider this time a lower limit for enrichment of the ISM since redistribution and cooling of the gas will occur on timescales of order  $10^8$  yr. This implies that the ISM enrichment process in J1148+5251 presumably started at redshifts  $z > 8$ . This is in agreement with the recent results from the Wilkinson Microwave Anisotropy Probe (WMAP) which indicate that the first onset of star formation most likely occurred at redshifts  $z \sim 15 - 17$  ( $\sim 250$  Myr after the Big Bang)<sup>25,26,27</sup>.

The presence of Ly $\alpha$  emitting galaxies and luminous quasars at the end of cosmic reionization ( $z > 6.3$ ), at a time when the IGM was at least 1% neutral, was clearly demonstrated<sup>5,28</sup>. This epoch of reionization represents a key benchmark in cosmic structure formation, indicating the formation of the first luminous structures. Detecting a large reservoir of molecular gas in this epoch demonstrates the existence of the requisite fuel for active star formation in primeval galaxies. The existence of such reservoirs of molecular gas at early times implies

that studies of the youngest galaxies will be possible at millimetre and centimetre wavelengths, unhindered by obscuration by the neutral IGM.

## References

1. Ohta, K. et al. Detection of molecular gas in the quasar BR1202-0725 at redshift  $z=4.69$ , *Nature* **382**, 426–428 (1996)
2. Omont, A. et al. Molecular gas and dust around a radio-quiet quasar at redshift 4.69, *Nature* **382**, 428–431 (1996)
3. Carilli, C.L. et al. High-Resolution Imaging of Molecular Line Emission from High-Redshift QSOs, *Astron. J.* **123**, 1838–1846 (2002)
4. Carilli, C.L. & Blain, A.W. Centimeter searches for molecular line emission from high-redshift galaxies, *Astrophys. J.* **569**, 605–610 (2002)
5. Fan, X. et al. A survey of  $z > 5.7$  quasars in the sloan digital sky survey. II. Discovery of three additional quasars at  $z > 6$ , *Astron. J.* **125**, 1649–1659 (2003)
6. White, R.L., Becker, R.H., Fan, X. & Strauss, M.A. Probing the ionization state of the universe at  $z>6$ , *Astron. J.* in press (2003)
7. Willott, C.J., McLure, R.J. & Jarvis, M.J. A  $3 \times 10^9 M_{\odot}$  black hole in the quasar SDSS J1148+5251 at  $z=6.41$ , *Astrophys. J.* **587**, L15–L18 (2003)
8. Loeb, A. & Barkana, R. The Reionization of the Universe by the First Stars and Quasars, *Ann. Rev. Astron. Astrophys.* **39**, 19–66 (2000)
9. Bertoldi, F., Carilli, C.L., Cox, P., Fan, X., Strauss, M.A., Beelen, A., Omont, A. & Zylka, R. Dust emission from the most distant quasars, *Astron. Astrophys. Letter* in press (2003)
10. Bertoldi, F. et al., *Astron. Astroph. Letters*, 2003, subm.

11. Solomon, P.M., Radford, S.J.E. & Downes, D. Molecular gas content of the primeval galaxy IRAS 10214+4724, *Nature* **356**, 318–321 (1992)
12. Strong, A.W., et al. Diffuse continuum gamma rays from the Galaxy observed by COMPTEL, *Astron. Astroph.*, **292**, 82–91 (1994)
13. Downes D. & Solomon, P. Rotating Nuclear Rings and Extreme Starbursts in Ultraluminous Galaxies, *Astroph. J.* **507**, 615–654 (1998)
14. Weiss, A., Neininger, N., Huttemeister, S. & Klein, U. The effect of violent star formation on the state of the molecular gas in M82, *Astron. Astrophys.* **365**, 571–587 (2001)
15. Solomon, P.M., Downes, D., Radford, S.J.E. & Barrett, J.W. The Molecular Interstellar Medium in Ultraluminous Infrared Galaxies, *Astroph. J.* **478**, 144 (1997)
16. Ferrarese, L. & Merritt, D. A fundamental relation between supermassive black holes and their host galaxies, *Astroph. J.* **539**, L9–L12 (2000)
17. Gebhardt, K. et al. A relationship between the nuclear black hole mass and galaxy velocity dispersion, *Astroph. J.* **539**, L13–16 (2000)
18. Richards, G.T. et al. Broad emission-line shifts in quasars: an orientation measure for radio-quiet quasars? *Astron. J.* **124**, 1–17 (2002)
19. Haiman, Z. & Cen, R. A constraint on the gravitational lensing magnification and the age of the redshift  $z=6.28$  quasar SDSS 1030+0524, *Astroph. J.* **578**, 702–707 (2001)
20. Elvis, M. et al. Atlas of quasar energy distributions, *Astroph. J. Sup.* **95**, 1–68 (1995)
21. Telfer, R.C., Zheng, W., Kriss, G.A. & Davidsen, A.F. The Rest-Frame Extreme-Ultraviolet Spectral Properties of Quasi-stellar Objects, *Astroph. J.* **565**, 773–785 (2002)
22. Pentericci, L. et al. VLT Optical and Near-Infrared Observations of the  $z = 6.28$  Quasar SDSS J1030+0524, *Astron. J.* **123**, 2151–2158



23. Heger, A. & Woosley, S.E. The nucleosynthetic signature of population III, *Astroph. J.* **567**, 532–543 (2002)
24. Arnett, A. Massive star evolution and SN 1987A, *Astroph. J.* **383**, 295–307 (1991)
25. Cen, R. Implications of WMAP Observations On the Population III Star Formation Processes, *Astroph. J. Letters* subm.
26. Kogut et al. Wilkinson Microwave Anisotropy Probe (WMAP) First Year Observations: TE Polarization, *Astroph. J.*, subm. (2003)
27. Spergel et al. First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters, *Astroph. J.*, subm. (2003)
28. Hu, E.M. et al. A redshift  $z=6.56$  galaxy behind the cluster Abell 370, *Astroph. J.* **568**, L75–L79 (2002)

**Acknowledgments:** The VLA is operated by the National Radio Astronomy Observatory (NRAO), a facility of the National Science Foundation (NSF), operated under cooperative agreement by Associated Universities, Inc. (AUI). This work is partly based on observations carried out with the IRAM Plateau de Bure Interferometer. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain).

**Competing Interest Statement:** The authors declare that they have no competing financial interests.

**Correspondence** should be addressed to F.W. (e-mail: fwalter@nrao.edu)

*Fig. 1: CO detection in J1148+5251. Top:* VLA CO(3–2) emission detected at 46.6149 GHz ( $z=6.418$ , bandwidth: 50 MHz,  $\Delta z=0.008$ ). *Bottom:* Co-added VLA emission in the neighboring line-free channels. The ( $1\sigma$ ) noise in both images is  $0.057 \text{ mJy beam}^{-1}$  and contours are plotted at -0.1, 0.1, 0.2, 0.3, 0.4 and  $0.5 \text{ mJy beam}^{-1}$ . The peak flux in the top panel is  $0.570 \text{ mJy beam}^{-1}$ ; i.e. J1148+5251 is detected at  $10\sigma$ . Based on the line free channels we derive a  $2\sigma$  upper limit to the continuum emission at 46.6 GHz (restframe wavelength= $870\mu\text{m}$ ) of  $0.10 \text{ mJy}$ . The cross indicates the optical position of J1148+5251. The optical and radio positions are coincident within the uncertainties ( $\sim 0.1'' = 0.6 \text{ kpc}$ ). The observations were taken with the VLA in the most compact configuration (D array, maximum baseline:  $\sim 1 \text{ km}$ , leading to a resolution of  $1.8'' \times 1.5''$ ; the beam is plotted in the lower left of each panel). Only two 50 MHz channels can be observed at once with the VLA, i.e., the source has been observed repeatedly with different frequency settings; observations were made using two polarizations and 2 IFs of 50 MHz each, scanning in frequency from 47.065 GHz to 46.515 GHz, corresponding to a CO(3–2) redshift range of 6.347 to 6.434. The observing time for the channels shown are  $\sim 20$  hours, respectively. Observations were obtained in fast switching mode, and the phase stability was excellent at all times. We used the nearby quasar 11534+49311 (flux density =  $1.6 \text{ Jy}$  at 46.6 GHz) for phase and amplitude calibration. The absolute flux calibration was derived by observing 3C286. No evidence for strong gravitational lensing is seen in optical images<sup>5</sup>, rendering strong magnification unlikely. However, the presence of an intervening galaxy at  $z=4.9$  has been suggested based on optical spectroscopy<sup>6</sup>, and counts of radio and optical sources in the vicinity of J1148+5251 argue for a foreground cluster, such that weak magnification ( $\sim$  factor two) is possible<sup>9,5</sup>.

*Fig. 2: The CO spectrum of J1148+5251.* The spectrum of the CO(3–2) line of J1148+5251 at  $320 \text{ km s}^{-1}$  resolution as observed with the VLA (top panel). The plotted errors are  $\pm 1\sigma$ . The averaged CO(6–5) and CO(7–6) observations from the PdBI are shown in the bottom panel<sup>10</sup> (channel width:  $5 \text{ MHz} = 13.8 \text{ km s}^{-1}$ , noise per channel:  $0.8 \text{ mJy}$ , beam size:  $\sim 5''$ ) to demonstrate the consistency of the results obtained by the two instruments. A Gauss fit to the PdBI data gives a velocity width of  $250 \text{ km s}^{-1}$  and a redshift of  $z=6.419$  (corresponding to ‘0’ velocity).



